

J<sup>r</sup> Loze 28

Papd March 10 1829

Dissertation  
on  
Respiration,  
by  
Amos Pennickaker  
of  
Pennsylvania.

22 Oct 2

Left at 10 AM

10 miles

10 miles

10 miles

10 miles

Pro

With

To  
William Gibson, M.D.  
Professor of Surgery in the University  
of  
Pennsylvania,  
this  
Dissertation  
is  
Dedicated,

With sentiments of the highest  
Respect and Esteem,  
By  
Amos Pencilaker.

The

because  
through  
profits  
return  
of the  
before  
is neg  
contact  
undergo  
deprive  
of carb  
fulcut  
consist  
The  
which  
upward  
res; the



## Respiration.

The blood, after having been sent by the left ventricle of the heart through the arteries to every part of the body, for its nourishment and support, is again returned, by the veins, to the right auricle of the heart, and supplied with chyle; but, before being distributed over the system, it is necessary that it should be brought in contact with the atmosphere, that it may undergo certain changes, by which it is deprived of its superabundant quantity of carbon; for this purpose there is a complicated apparatus called respiratory, consisting of the larynx, trachea, and lungs.

The larynx is composed of five cartilages, which form an irregular tube that opens upwards, corresponding to the posterior nares; the inferior portion is connected to

the  
of the  
holl  
flat  
his  
two  
anto  
it ad  
by an  
is al  
rings  
When  
it in  
the pe  
into t  
verte  
the ly  
suble  
-w in

the trachea, which is placed in the fore part of the neck, anterior to the oesophagus. It is a hollow tube, cylindrical in front, but flat on its back part, consisting of a number of cartilaginous rings that form about two thirds of the circumference of a circle anteriorly, the third behind, where there is a deficiency in the rings, is filled up by an elastic fibrous membrane which is also placed in the intervals between the rings and serves to connect them together. When the trachea passes into the thorax, it inclines a little backwards and enters the posterior mediastinum, and divides into two branches opposite the third dorsal vertebra, one going to the right, the other to the left lung. These branches are again subdivided into smaller ones which, together with the two main branches, are na-

-sued  
the to  
trach  
memb  
After  
-nes, to  
I com  
the bus  
-ped to  
ready  
-ber of  
in every  
appear  
having  
a cell,  
also to  
of the b  
atmosph  
in the

-and bronchiae. The internal surface of the bronchia, as well as the larynx and trachea are lined by a soft elastic mucous membrane.

After having mentioned in a cursory manner, the canal and its branches, by which a communication is established between the lungs and external air, it may be proper to describe the lungs. The bronchia already mentioned divide into a great number of branches which ramify in the lungs in every direction, and now appears the appearance of small membranous tubes, having attached to their extremities vesicles or cells, which differ from the common cellular texture of the lungs and other parts of the body. It is in these cells that the atmosphere effects the necessary changes in the blood which is conveyed to them

by the  
 take  
 triole  
 sides  
 rest of  
 left  
 so of  
 which  
 such  
 around  
 with  
 with  
 large  
 they  
 the  
 been  
 delicate  
 Base  
 on the

by the pulmonary arteries. These arteries take their origin at the base of the right ventricle by 2 common trunks, that soon divide into two branches, one entering at the root of the right lung, the other at that of the left; when they again divide and subdivide, so as to become small capillary vessels, which ramify around the air cells in such a manner as to form a net work around them, and then they interminate with a set of capillary veins, these unite with one another, and at length form two large trunks for each lung, from which they emerge to discharge their blood into the left auricle of the heart, after having been subjected to the atmosphere in the delicate vessels around the air cells.

Besides the pulmonary arteries there are others for the nourishment of the lungs

ordin  
ordres  
of the  
the br  
belong  
into the  
long  
The  
numb  
their  
their  
in the  
which  
around  
lungs  
with  
from  
These  
er by



arising from the aorta, these are the bronchial arteries which penetrate the different tissues of these viscera, and finally terminate in the bronchial veins; those of the veins which belong to the right lung discharge their blood into the venæ cavae, and those which belong to the left, into the subclavian vein. The lungs are also furnished with a great number of absorbent vessels that arise from their internal substance, as well as from their external surface. They are directed in their course towards the bronchial glands, which are of a blackish colour and placed around the bronchies where they enter the lungs. Lastly, the lungs are supplied with nerves which are principally derived from the par vagum and intercostals.

These different tissues, are united together by an intermediate cellular tissue,

which  
gettle  
by of  
each  
The  
symp  
of two  
conci  
lobes)  
and A  
sed b  
havin  
impro  
to live  
pulsar  
the sac  
is nam  
The d  
oid, to

which not only serves to connect them together; but is also useful in keeping the vessels of these organs from encroaching upon each other.

The lungs thus constituted of arteries, veins, lymphatics, &c. &c. present the appearance of two large spongy bodies of an irregular coincident shape, each one, is divided into lobes by fissures, the left consisting of two, and the right of three; they are also inclosed by a serous membrane, each lung having one that may be compared to an imperious sack, one side of which serves to line the lung and is designated pleura pulmonalis, and that side or portion of the sack which lines the cavity of the chest is named pleura costalis.

The thorax, in which the lungs are placed, has in shape some resemblance to

a con  
base  
by the  
by the  
each  
partic  
super  
the st  
angue  
-cuse  
and  
at wa  
and  
and  
also  
-gm,  
of the  
twice  
these

a cone, the apex of which is above, and the base below; the posterior portion is formed by the dorsal vertebrae, and its lateral sides by twenty four ribs, twelve being placed on each side, and articulated behind with the vertebrae; before, the seven that form the superior section of the chest articulate with the sternum, a bone that is somewhat triangular in form; these bones form an inclosure for the protection of the heart and lungs, and have attached to them, at various places, muscles for increasing and diminishing the capacity of the chest, and a variety of other purposes. There is also a circular muscle called diaphragm, attached to the outline of the base of the chest, that forms a partition between the viscera of the thorax and those of the abdomen.

The  
with  
condens  
and p  
necessa  
the ma  
-form  
fluid  
worth  
of large  
dens,  
in pro  
-presin  
to dimi  
and op  
it will  
also the  
is unde  
six; su

The atmosphere is so intimately connected with the function of respiration, that a consideration of some of its ~~most~~ chemical and physical properties are essentially necessary to a perfect understanding of the manner in which venous blood is transformed into arterial. It is that aeriform fluid which every where surrounds the earth to the height of a considerable number of leagues; it is invisible, colourless, without odour, and only sensible to the touch when in motion; it has the property of compressibility; for a portion may be made to diminish greatly in volume by pressure, and as soon as the pressure is removed it will resume its former bulk, proving also that it possesses elasticity. There is another property in atmospheric air, namely its weight, which may be

use  
cause  
then  
will  
the  
side  
it  
this  
from  
be  
are  
greater  
consider  
it is  
are  
these  
ating  
have  
the



ascertained by exhausting a vessel, and counterpoising it on a scale beam, and then by allowing the air to rush in, it will be found that the end to which the vessel is attached will preponderate considerably. It is owing to its gravity that it presses equally on every part of the earth's surface; its pressure is about fifteen pounds to every square inch. It must be obvious that the strata of air which are near the earth's surface sustain a greater weight than those that are a considerable distance from it, and as it is compressible, those portions which are below must be more dense than those which are more elevated. These variations in the density of the atmosphere have no influence on the functions of the human body when near the earth,

but a  
- sub  
size  
and  
in the  
thing  
in the  
of ex  
of m  
hous  
- sub  
prop  
labor  
- ce p  
These  
nec  
for if  
wou  
rem

but when man ascends to the tops of high mountains, where the pressure of the air is not so great, and consequently it must be lighter and rarer, a remarkable effect is produced in some of the animal and mental functions; owing to the diminished pressure on the surface of the body and deficiency of oxygen, there is a sudden exhaustion of muscular power, palpitations of the heart, throbbing of the arteries, universal weakness, accompanied with a propensity to sleep, respiration becoming laborious; lastly, hemorrhage takes place from various parts of the body.

These derangements serve to show the necessity of the atmospheric pressure; for if its pressure were removed, it would be impossible for our bodies to remain in the state in which we see

thou  
thou  
-say  
were  
-nat  
by li  
I  
prop  
prose  
if cry  
adver  
they a  
of cry  
ding  
-ing  
-lity  
that  
ent  
agress

them, the fluids would be determined to their external surfaces, and thereby derange the internal functions; hemorrhage would also take place from ~~various~~ internal parts that are but slightly protected by linings like the mucous membranes.

I shall now proceed to the chemical properties of the atmosphere. It was supposed to be an element, till the discovery of oxygen, when its real composition was ascertained by Scheele and Lavoisier, though they differed with regard to the quantity of oxygen it contained. Lavoisier, according to his analysis, fixed it at 27 ~~measures~~ <sup>measures</sup>; Scheele supposed the quantity to be still greater. Subsequent to that time, it has been analysed by different chemists with more precaution, and, agreeably to the best authority, the properties

of the  
about  
77.5  
any  
acc  
is a  
-ous  
that  
be a  
just  
the  
a co  
and  
other  
than  
-es:  
yet  
-duc  
was

of the different gases are represented to be about 26 of oxygen by measure, nitrogen 77.8; Besides these gases, there enter into the composition of the atmosphere some carbonic acid and moisture; the quantity of carbonic acid is, by measure, 0.03, and that of aqueous vapour 142; these are the constituents that compose the atmosphere, and it must be evident from the analysis that has just been stated, that nitrogen forms the greater part of its bulk, which is a colourless gas destitute of taste or odour, and may be better distinguished from other gases by its negative properties than any peculiar character it possesses: it does not support life or combustion, yet when inhaled into the lungs, it produces no injurious effects on them. It was at one time supposed that its use

was  
it is  
imp  
-gu  
-im  
en  
and  
oxy  
than  
with  
is c  
or lo  
supp  
ent  
m  
ave  
depe  
life  
in a



was merely to dilute the oxygen, but it is more probable that it may serve some important purposes of which we are unacquainted, for it has been proved by experiment, that respiration can be carried on without its being mixed with oxygen and that when an animal respirs pure oxygen, there is no more of life consumed than if it were used in a state of mixture with nitrogen. Oxygen, like nitrogen, is colourless, transparent, without odour or taste, but possesses the property of supporting life and combustion, and enters into all animal and vegetable matter. It is this gas to which the air owes its chemical properties, for, when deprived of it, it will cease to support life or combustion; metals when heated in it will no longer become oxidized,

of  
neg  
me  
the  
the  
py  
its  
form  
it  
const  
It is a  
and  
tion  
there  
her  
time  
dual  
-  
time

as when oxygen is present. When the necessity of air is experienced, a number of muscles are called into action, by which the capacity of the chest is increased; the air, as has already been mentioned, presses on every part of the body, owing to its gravity, therefore, when a vacuum is formed in the lungs and their canal, it will consequently rush into them: this constitutes the process denominated inspiration. It is again expelled or expired by the action of another set of muscles, which by their contraction diminish the capacity of the chest, and thereby produce the process named expiration. The number of inspirations in a given time vary considerably in different individuals, and also in the same person, ~~for~~ <sup>for</sup> a man that respires twenty times in a minute at one time, may at

an  
so of  
by a  
step  
tion  
nect  
-wre  
air to  
tion.  
diff  
ing  
good  
incl  
2, ut  
to be  
that  
that  
lungs  
cairle

another respires more frequently or not so often; the number is greatly influenced by a variety of circumstances, such as sleep, violent exercise, and moral afflictions. There is another circumstance connected with inspiration that is not unworthy of notice, namely, the quantity of air that enters the lungs at each inspiration. Various estimates have been made by different physiologists, but without coming to a uniform conclusion; thus by Goodwin, it has been stated to be 12 cubic inches, by Buvier 18 1/4, and by Gregory 2, while Doctor Thompson thinks it to be 40 cubic inches; he also thinks that the same amount is expired, and that there is generally about 250 in the lungs. The air, by traversing the nasal cavity, mouth, larynx, trachea and

bro  
se as  
it was  
express  
by the  
-dys  
napre  
mem  
Alen  
in wh  
lung  
-dys  
to not  
chang  
lung  
tion  
-alut  
probab  
been in

bronchia has its temperature increased  
so as to be equal with that of the body;  
it must therefore become rarer and  
expanded, so as to increase in volume  
by the time that it enters the lungs, besi-  
des these changes it becomes mixed with  
vapour as it passes along the mucous  
membrane of the air passages.

After having mentioned the manner  
in which the air gains access to the  
lungs, and the slight changes it un-  
dergoes, it will be my next object  
to notice the chemical and physical  
changes it has undergone while in the  
lungs: In a few seconds after its introduc-  
tion into these organs, a quantity, equiv-  
alent to that inspired, is expired, though  
probably not the same that has just  
been introduced in the preceding inspi-

- you  
 it is  
 - you  
 the  
 been  
 - me  
 - for  
 but  
 - you  
 for a  
 so of  
 that  
 when  
 only  
 un-  
 red  
 must  
 long  
 air's



-ration, however, whether it be the same  
or not, it is very evident that it has under-  
gone certain alterations, for instead of  
the same quantity of oxygen that had  
been introduced, we find it greatly di-  
minished, while the nitrogen has suf-  
fered neither increase nor diminution in  
bulk. The carbonic acid has also under-  
gone a change with regard to quantity,  
for it will be found to have increased  
so as to represent the quantity of oxygen  
that has disappeared, which was 0.21  
when inhaled, but now there remains  
only 0.18, & 0.19, while the carbonic acid  
amounts to 0.2 & 0.21. The air that is expi-  
red also is mixed with vapour, which  
must probably exhalate from the capil-  
lary vessels that ramify around the  
air cells. It has been supposed that

the  
the  
who  
plea  
-has  
put  
-sue  
The  
order  
the  
lung  
has  
-fice  
prop  
ring  
my  
the  
-ing  
at the

the pulmonary exhalation is equal to the cutaneous, the extent of surface from which pulmonary exhalation takes place, and the fact that cutaneous exhalation is in an inverse ratio with the pulmonary, seem to warrant the conclusion.

The transformation of venous blood into arterial - the venous blood is conveyed from the right ventricle of the heart to the lungs by the pulmonary arteries, and has to pass through their minute ramifications around the air cells; it then passes into the radicals of the pulmonary veins; while thus traversing these capillary vessels it is brought in contact with the atmosphere in the air cells, and by parting with its carbon is converted into arterial blood, which will be found to be of a

sew  
ed  
from  
er to  
more  
greater  
sown  
the ac  
in the  
shall  
take p  
that t  
the air  
vapour  
cupped  
and w  
dimin  
respir  
attract

scarlet red colour, and to have a stronger  
colour than the venous, differing likewise  
from the venous in being possessed of a high-  
er temperature, and the property of a  
more rapid coagulation; it has also a  
greater specific gravity, and contains less  
serum. The changes that take place in  
the air and blood, when brought together  
in the lungs, having been described, I  
shall next endeavour to show <sup>how</sup> some of them  
take place. It has already been mentioned  
that there is a portion of vapour mixed with  
the air that is expired; the formation of this  
vapour is owing to an exhalation from the  
capillary branches of the pulmonary arteries,  
and will most satisfactorily account for the  
diminution of serum in the blood during  
respiration. The change of colour has been  
attributed to iron in the blood, but it is

...  
with  
the  
-gon  
-co,  
phys  
cart  
lary  
at the  
cart  
that  
throu  
the c  
and  
Bisio  
there  
-sides  
resp  
of there  
its, av

more probable that this alteration, as  
well as its increase of amount, depends upon  
the disengagement of carbon from its re-  
-gards, the manner in which this takes pla-  
-ce, various opinions are entertained by  
physiologists; some supposing that the  
carbon is transmitted through the capil-  
lary parities into the air cells, and that  
it there combines with the oxygen, forming  
carbonic acid, while others again suppose  
that the oxygen is absorbed by the blood  
through the capillary vessels, and that  
the carbonic acid is formed in them  
and afterwards passes into the cells.  
Besides the chemical actions mentioned,  
there are vital ones which have con-  
-siderable influence on the blood during  
respiration; it is owing to the conjoint agency  
of these two actions that the blood changes  
its various character in the lungs. A P